

**GEOLOGIC MAP OF THE
OSIRIS (Jg-12)
AND
APSU SULCI (Jg-13)
QUADRANGLES
OF GANYMEDE**

By Don E. Wilhelms

1997

Prepared for the National Aeronautics and Space Administration
by
U.S. Department of the Interior, U.S. Geological Survey

(Published in hardcopy as USGS Miscellaneous Investigations Series Map I-2442, as part of the Atlas of Jovian Satellites, 1:5,000,000 Geologic Series. Hardcopy is available for sale from U.S. Geological Survey, Information Services, Box 25286, Federal Center, Denver, CO 80225)

Please direct questions or comments about the digital version to:

Richard Kozak
U.S. Geological Survey
2255 N. Gemini Drive
Flagstaff, AZ 86001

e-mail: rkozak@flagmail.wr.usgs.gov

DESCRIPTION OF MAP UNITS

LIGHT MATERIALS

- s** **Smooth material**—Forms predominantly smooth or slightly hummocky surfaces. Delimited by solid contact lines or structure symbols where sharply bounded; delimited by dashed contact lines where apparently gradational with grooved material. Queried where obscured by rays and could be grooved, palimpsest, or dark material. *Interpretation:* Undeformed or slightly deformed ice derived from extruded water or slush. Patches distal to Gilgamesh Basin may be Gilgamesh ejecta
- g** **Grooved material**—Forms domains predominantly consisting of two or more parallel grooves, each 3–10 km wide and as long as 500 km; most grooves gently curved but some straight or sharply curved; raised rims and U-shaped profiles revealed by low-angle lighting. Queried where obscured by rays and could be dark or reticulate material. *Interpretation:* Ice derived from extruded water or slush like that of smooth material, but deformed after extrusion
- gf** **Grooved material, fine**—Forms generally small domains. Grooves and ridges short and narrow, commonly sinuous, produce finer overall texture than in grooved unit but coarser than in smooth unit. *Interpretation:* Materials like those of grooved unit but less severely deformed, possibly because thinner
- l** **Light material, undivided**—Albedo typical of light units (0.3–0.5) but topographic features too poorly imaged for further identification; queried where obscured by rays and could be dark material. *Interpretation:* Ice derived from extruded water or slush

RETICULATE MATERIALS

- r** **Reticulate material**—Cut by short grooves meeting at high angles. Intermediate in albedo between typical dark and light materials, commonly darkest in the grooves, brightest on the highs. *Interpretation:* Dark material disrupted by gridlike faults and coated by thin ice. Predates nearby grooved material
- dr** **Dark reticulate material**—Similar to reticulate material (unit r) but darker. *Interpretation:* Formed early in breakup of dark crust; has little coating of fresh ice

DARK MATERIALS

- ds** **Dark smooth material**—Forms relatively smooth surfaces, but more highly textured than brighter smooth material (unit s) except in large, smooth, lobate patch centered at lat 32° S., long 186°. *Interpretation:* Extruded ice containing some silicate material. Patches distal to Gilgamesh Basin may be Gilgamesh ejecta
- dl** **Dark lineated material**—Contains linear grooves more widely spaced and with rougher edges than grooves of light grooved material. Queried where obscured by rays and could be reticulate or light grooved material. *Interpretation:* Structurally deformed dark material, possibly precursor of light grooved material
- dc** **Dark cratered material**—Contains many degraded craters and crater segments. Darkest extensive unit in map area. *Interpretation:* Less modified than other units but probably younger than original crust
- df** **Dark furrowed material**—Extended surfaces with widely spaced furrows a few kilometers wide having raised, scalloped, ragged rims. Greatest furrow concentration in map area forms subcircular system concentric around point in

- palimpsest at lat 28° S., long 153°. *Interpretation:* Ancient crust fractured by shock waves from large impacts followed by structural modification that opened the fractures (Schenk and McKinnon, 1987)
- dh **Dark hummocky material**—Forms small elevated patches having rough surfaces. *Interpretation:* Ancient ejecta or intensely deformed part of ancient crust
- d **Dark material, undivided**—Albedo typical of dark units (0.2–0.4) but topographic features too poorly imaged or too limited in extent for further identification. Queried where obscured by rays and could be a light or reticulate unit

CRATER AND PALIMPSEST MATERIALS

[Only craters 20 km in diameter and larger are mapped]

- c3 **Crater material, unit 3**—Interior and exterior materials of sharply textured, bright-rayed craters. Some albedos as high as 0.73 (McKinnon and Parmentier, 1986). *Interpretation:* Youngest impact craters in the map area and probably younger than all other units
- cs3 **Secondary crater material, unit 3**—Interior and exterior materials of small, clustered craters within or near crater rays. (Rays not separately mapped because most show on base map.) *Interpretation:* Secondary-impact craters responsible for excavating dark and light substrates
- c2 **Crater material, unit 2**—Interior and exterior materials of well-formed, rayless craters. *Interpretation:* Impact craters of intermediate age formed throughout long time period
- cs2 **Secondary crater material, unit 2**—Interior and exterior materials of small, clustered craters adjacent to rayless craters. *Interpretation:* Secondary-impact craters and their deposits
- c1 **Crater material, unit 1**—Interior and rim materials of highly degraded craters. *Interpretation:* Either old or anomalously degraded
- cd **Crater material, dark**—Forms floors and patches on flanks of some craters, including those too small to map at lat 22.5° S., long 141°; lat 27.5°, long 167.5°; and lat 28.5° S., long 194°. Albedo as low as 0.11 (Schenk and McKinnon, 1991). *Interpretation:* Ice contaminated by projectile, or extrusions of material contaminated by silicates
- p **Palimpsest material**—Forms circular or elliptical patches intermediate in albedo and smoothness between surrounding dark and light materials. Queried where no rounded form seen. *Interpretation:* Material of old craters or basins flattened by relaxation of crust; queried occurrences could be extruded thin light material


GILGAMESH BASIN MATERIALS

- bf **Basin floor material**—Smooth to slightly textured material composing central depressed floor of Gilgamesh. *Interpretation:* Either melted target material or later interior deposit
- br **Basin rugged material**—Coarsely hilly interior and rim materials of Gilgamesh. May form outlying patches subconcentric with basin and containing subconcentric small ridges and hummocks. *Interpretation:* Primary ejecta of Gilgamesh and deformed target material. Outlying patches possibly decelerated ground-flow ejecta
- bs **Basin smooth material**—Smooth or gently hummocky material transitional with




















rugged material. *Interpretation:* Gilgamesh primary ejecta, possibly mixed with some locally derived substrate material

csb **Secondary crater material of basin**—Interior and surrounding deposits of clustered and elongate craters as much as 20 km across, arrayed radially to Gilgamesh. *Interpretation:* Secondary-impact craters of Gilgamesh and their ejecta

CORRELATION OF MAP UNITS



*Age range includes dark smooth material (unit ds)

	Contact —Dashed where gradational; dotted where buried; queried where obscured by rays (especially west and south of Osiris), by shadows (especially near terminator), or by poor photographic resolution. Includes groove domain boundaries
	Long, linear trough —Dotted where subdued; queried where continuity uncertain. Similar to one or pair of grooves in light grooved material, but generally longer and deeper; may extend into dark terrain; includes flat bottomed, narrow troughs. Interpreted as master fracture along which dark terrain broke up most deeply. Apparently buried where subdued
	Deep, short linear trough in dark and light materials —Interpreted as relatively young fracture
	Sharp groove trend —In grooved material; schematic
	Subdued groove trend —In grooved or smooth material; schematic
	Reticulate groove trend —Schematic
	Furrow in dark and palimpsest materials —Narrow, linear depression with raised, scalloped rim; shorter and more irregular than long linear troughs and more widely spaced than light-material grooves. Interpreted as graben formed by modification of impact-related arcuate fractures (Schenk and McKinnon, 1987)
	Lineament in dark material —Schematic
	Ridge in light material —Dotted where buried. May be caused by near surface compression
	Sinuuous rille —Narrow, curvilinear depression. Possibly formed by flowing water
	Irregular depression —Widening at end of sinuous rille. Possibly site of water eruption
	Crater rim crest
	Buried crater rim crest
	Inward-facing scarp on crater floor —Line marks top
	Peak on crater floor
	Pit on crater floor —Dot where too small to map
	Dome on crater floor —Line marks base
	Palimpsest ring
	Basin ring arc

GENERAL FEATURES

A mosaic of dark-toned and light-toned geologic units about equal in total area characterizes Ganymede, whose diameter of about 5,262 km makes it Jupiter's and the Solar System's largest satellite (Burns, 1986). Ganymede's low bulk density (1.94 g/cm^3 ; Burns, 1986), spectral properties, and high albedos indicate that water ice is the main component of both the dark and the light units of the crust (Smith and others, 1979; Johnson and others, 1983; Clark and others, 1986; McKinnon and Parmentier, 1986). Impact craters pepper but do not saturate the surface.

Light units predominate in the map area, which includes the parts of two quadrangles that were imaged at geologically useful scales and illuminations by Voyager 2 during its flyby of the Jovian system in July 1979 (Smith and others, 1979). Ganymede is locked in synchronous rotation so that longitudes 90° – 270° , including the map area, always face away from Jupiter. Territory east of long 180° , including the Osiris quadrangle (Jg-12), is on the hemisphere that leads in Ganymede's revolution around Jupiter; territory west of 180° , including the Apsu Sulci quadrangle (Jg-13), is on the trailing hemisphere. The southeastern corner of the map area includes Ganymede's and the Jovian system's largest impact structure with well-preserved morphology, the Gilgamesh Basin.

DARK AND LIGHT MATERIALS

A somewhat higher crater density of the dark materials (Shoemaker and others, 1982; Chapman and McKinnon, 1986; Murchie and others, 1989) and interruption of the dark units and their structures by lanes, wedges, and polygons of light materials demonstrate that the dark are mostly, though not wholly, older than the light materials (Lucchitta, 1980; Golombek and Allison, 1981; McKinnon and Parmentier, 1986; Murchie and others, 1986; Squyres and Croft, 1986). Contamination of pure ice by less than 10 and possibly only 3 percent of dark grains, possibly of cometary and meteoroid origin, can explain the albedo difference between the dark (0.2–0.4) and light (0.3–0.5) materials (Clark and others, 1986; McKinnon and Parmentier, 1986). The generally older dark units have had more time to accumulate this debris. They are subdivided on the basis of superposed craters or structures such as furrows and lineations. Except for the dark smooth material (unit ds), all dark units have been modified since formation, and none is believed to represent the original crust (Murchie and others, 1989, 1990). Embayment of older units and structures suggests that at least one patch of dark smooth material (centered at lat 32°

cratered material at lat 62° S., long 156°). Only minor lateral shear is indicated in the map area, an interpretation also made for the whole of Ganymede by Schenk and McKinnon (1987); however, Murchie and Head (1988) favored major shear and rotation in many regions.

Ganymede's principal tectonic regime seems, instead, to have been crustal extension on the order of 5 to 7 percent caused by global expansion on the order of only 1 percent (Squyres, 1980; McKinnon, 1981; Golombek, 1982; Squyres and Croft, 1986). Emplacement of light materials therefore required less profound modification of the crust than might be implied by their great extent. The crust stretched enough to open long, linear troughs (bowtie symbol) and short, ragged lineations in patches of dark lineated material (unit dl) and reticulate materials. Some of the linear troughs cross only dark materials, but most of them or their subdued, probably thinly buried extensions (for example, those near lat 32.5° S., long 129°) bound patches of reticulate or light materials. The fracturing allowed small amounts of water or slush to flow or spray out upon the surface. Most light units originated when larger amounts of this new material filled shallow depressions created where blocks of dark material dropped passively along the visible and buried faults (McKinnon and Parmentier, 1986; Squyres and Croft, 1986). Elevation differences consistent with the downdrop of dark crustal materials are observed along many dark-light contacts in the favorably illuminated northeastern part of the map area.

The most typical light material, the grooved unit (unit g), is characterized by narrow, regularly spaced, parallel grooves bundled in sets or domains of two or more grooves and locally measuring as much as 100 km in width. Some groove sets parallel long linear troughs, forming what Murchie and others (1986) called groove lanes. Other groove sets terminate against the troughs or domains of light material in a T-relation, in which the crosscutting structure is commonly the older (Golombek and Allison, 1981; Murchie and Head, 1988). The most common orientation of the troughs and largest grooves in the map area is west-northwest. In the northeastern part of the map area, other, seemingly random trends lie between the west-northwest structures. Some major troughs and grooves veer off to the west-southwest at about long 180° and, in the Apsu Sulci quadrangle, conspicuously truncate the west-northwest structures.

Light material that is smooth on most available images (unit s) is also abundant. It constitutes some entire domains, but more commonly it forms parts of domains in which grooved material is the other component. A few craters superposed on the smooth material are cut by grooves (lat 24° S., long 139°–141°). Therefore, the processes that formed grooves and light materials, although related, are not identical (Squyres and Croft, 1986). Some patches of smooth material embay grooved and other older units in the manner of an extruded fluid (for example, at lat 25.5° S., long 126.5°; lat 43.5° S., long 128°). Smooth units and groove sets intergrade, however, in most domains of light material (for example, at lat 22° S., long 151°; lat 31° S., long 130°), as shown by dashed contacts or suggested by different densities of schematic groove symbols. In these mixed domains, the smooth material apparently was the first to form, followed by partial grooving. A reasonable inference is that domains consisting entirely of grooved material also began as smooth material.

The global expansion that fractured the dark crust and allowed the light materials to rise and reach the surface may have directly caused some grooving as well, at least in the groove lanes (Murchie and others, 1986). A more important determinant of local groove structure may be the familiar property of the ordinary low-pressure phase of water ice

middle froze and expanded. The finely grooved material (unit **gf**), which forms relatively small domains and lacks the many parallel, deep, raised-rim grooves of the grooved material, may consist of thinner ice that simply froze, expanded, and deformed itself within its confines without arching. On the map, each domain that is believed to have been mechanically isolated during groove formation by any mechanism is bounded by fault symbols or solid contact lines.

CRATERS AND PALIMPSESTS

The impact origin of most craters on Ganymede is not disputed. Many clusters and fields of satellitic craters having subequal sizes (map units **CS2** and **CS3**, and unmapped where scattered) attest to the pervasiveness on Ganymede of the common Solar System process of secondary impact. Only a few small pits at the heads of apparent sinuous rilles show the signs of internal origin; flowing water emanating from these pits may have cut the rilles.

The age of a crater relative to adjacent geologic units is usually clear, but crater ages cannot easily be correlated in detail from region to region. Most nonrayed craters are therefore assigned to a single map unit (**C2**), and only a few highly degraded craters or crater fragments are assigned an older age (unit **C1**). As on other moons and planets, craters whose ejecta created rays during secondary impact are generally the youngest (unit **C3**). Rays obscure other geologic units west and southwest of Osiris, which has a diameter of 105 km and therefore, is the largest and most conspicuous young crater in the map area. Crater rays consist mainly of substrate material exposed and ejected by the secondary impact of the primary crater's ejecta. Ray brightness, therefore, depends partly on the brightness of the substrate unit: brightest on light materials and darkest on dark. The density of mapped craters (20 km and larger) is greater east of long 180° than west of it, an observation that fits the general finding that more craters of the mapped sizes were formed on the leading hemisphere than on the trailing hemisphere during a given time interval (Passey and Shoemaker, 1982). This distribution suggests that the impacting bodies revolved around the Sun, not Jupiter (Murchie and others, 1989).

The weakness of Ganymede's icy crust has greatly affected the appearance and retention of craters. The transition from craters having simple bowl-shaped interiors to those with central peaks occurs at crater diameters of only 3–6 km, well below the 19–27 km for the Moon (Passey and Shoemaker, 1982; Schenk, 1991). Therefore, all the craters mapped here are complex. Because surface gravity is similar on Ganymede and the Moon (144 and 162 cm/s², respectively), substrate properties must be a more important factor controlling the onset of the rebounds that create peaks than is gravity (Schenk, 1991). Schenk (1991) also cited crustal weakness at the time of impact as the cause of a relative shallowness of Ganymede's craters. Some craters larger than 25 km across possess central pits in or instead of the peaks, and craters larger than 50 km possess only pits (Schenk, 1991). The pits may originate by collapse of the peaks (Passey and Shoemaker, 1982) or by some as yet undetermined mechanism.

Apparently, the crust was especially weak early in its history because of a relatively high heat flow (McKinnon and Parmentier, 1986). One result is the flattening, probably by viscous relaxation (Passey and Shoemaker, 1982; Shoemaker and others, 1982), of old, large craters and basins into approximately circular, light-toned patches called palimpsests (Smith and others, 1979). Viscous relaxation or ice volcanism (Murchie and others, 1989, 1990) may also have destroyed many of the smaller old craters that must have formed along with the large ones. Later, cooling led to a more nearly elastic behavior of the crust and to the retention of younger craters (units **C2** and **C3**) (Hillgren and Melosh, 1989).

THE GILGAMESH BASIN

The Gilgamesh impact structure is considered a basin rather than a crater because of its

large size and multiple concentric, mountainous ring arcs. The relatively low density of superposed craters and the superposition of its deposits and secondary craters (unit *csb*) on almost all other nearby units show that it is relatively young, about 3.5 billion years, according to an interpretation of its crater density based on correlation with lunar crater densities (Shoemaker and others, 1982). The Gilgamesh Basin has not flattened out into a palimpsest and is not surrounded by concentric furrow systems like other basin-scale craters on Ganymede; therefore, the lithosphere had presumably cooled and thickened by the time it formed (Chapman and McKinnon, 1986). Concentration of the secondary craters north and south of Gilgamesh indicate that the impacting body approached approximately in the equatorial or ecliptic plane.

Gilgamesh is typical of lunar basins in some ways but atypical in others. Its many secondary craters, which appear in abundance about 500 km from its center, closely resemble those of the lunar Orientale Basin in size (as much as 20 km across), in shape (subcircular to elongate), and in their radial clustering (Wilhelms, 1987, chapter 4). Outward from its floor material (unit *bf*), the deposits of Gilgamesh grade from a rugged-appearing material (unit *br*) to a smoother facies (unit *bs*) that shows some evidence of outward flow in the form of indistinct lobes. Outlying patches of the rugged unit (quarried) also suggest ground flow, for they resemble basin-concentric ridges ("dunes") that were created around lunar basins when ejecta flowing along the surface piled up against obstacles.

However, the ring arcs of Gilgamesh are less well defined than those of most lunar basins (possibly another effect of the thick lithosphere; Chapman and McKinnon, 1986). Not even the longest and most conspicuous arc, whose radius averages about 275 km, defines an unambiguous topographic basin rim (the ring that encloses most of a basin's depression). The appearance of abundant secondary craters about 500 km from the center would indicate a basin diameter of about 500 km on the Moon (Wilhelms, 1987, chapter 4). However, this distribution of secondary craters suggests a diameter closer to 450 km on Ganymede because ejecta particles can fly farther under Ganymede's slightly lower surface gravity. Even well-defined topographic rims, however, may or may not bound the excavation cavities of lunar and planetary basins (Wilhelms, 1987, p. 77-81). At least it seems clear that Gilgamesh's excavation was not limited to its best-defined circular feature—the inner, smooth-floored depression 125 km in diameter—for the vastly simpler crater Osiris has a diameter only 20 km less in diameter (Lucchitta and Ferguson, 1988). Possibly Gilgamesh has no single well-defined excavation cavity, but rather a poorly defined zone in which excavation is greatest at the center and increasingly irregular and discontinuous farther out.

GEOLOGIC HISTORY


After accretion, Ganymede differentiated into upper layers rich in ice and lower layers and a core containing more silicate material (McKinnon and Parmentier, 1986). Repeated impacts created shallow craters and basins that relaxed even further into pancake-like palimpsests or that disappeared altogether. Dark cosmic material progressively darkened the surface. Dark ice volcanism may have resurfaced part or all of the ancient crust. Part of the crust was fractured by slight global expansion, possibly caused by the conversion of dense phases of water ice in the interior into less dense phases (Squyres and Croft, 1986). The dark terrain subsided slightly where intensely fractured, and water or slush from the interior filled the resulting lows. Global expansion and the freezing and expansion of the water or slush as ordinary ice created rimmed grooves over much, though not all, of its newly occupied area. Impacts continued throughout this history, and the resulting craters have remained moderately well preserved since the major disruption of dark materials and emplacement of light materials. Ganymede's largest impact into the thick lithosphere of this relatively late period created the Gilgamesh Basin about 3.5 billion years ago.

REFERENCES CITED



- Burns, J.A., 1986, Some background about satellites, *in* Burns, J.A., and Matthews, M.S., eds., *Satellites*: Tucson, University of Arizona Press, p. 1–38.
- Chapman, C.R., and McKinnon, W.B., 1986, Cratering of planetary satellites, *in* Burns, J.A., and Matthews, M.S., eds., *Satellites*: Tucson, University of Arizona Press, p. 492–580.
- Clark, R.N., Fanale, F.P., and Gaffey, M.J., 1986, Surface composition of natural satellites, *in* Burns, J.A., and Matthews, M.S., eds., *Satellites*: Tucson, University of Arizona Press, p. 437–491.
- Golombek, M.P., 1982, Constraints on the expansion of Ganymede and the thickness of the lithosphere, *in* Lunar and Planetary Science Conference, 13th, Houston, March 15–19, 1982, *Proceedings, Part 1: Journal of Geophysical Research*, v. 87, suppl., p. A77–A83.
- Golombek, M.P., and Allison, M.L., 1981, Sequential development of grooved terrain and polygons on Ganymede: *Geophysical Research Letters*, v. 8, p. 1139–1142.
- Hillgren, V.J., and Melosh, H.J., 1989, Crater relaxation on Ganymede: Implications for ice rheology: *Geophysical Research Letters*, v. 16, p. 1339–1342.
- Johnson, T.V., Soderblom, L.A., Mosher, J.A., Danielson, G.E., Cook, A.F., and Kupferman, Peter, 1983, Global multispectral mosaics of icy Galilean satellites: *Journal of Geophysical Research*, v. 88, p. 5789–5805.
- Lucchitta, B.K., 1980, Grooved terrain on Ganymede: *Icarus*, v. 44, p. 481–501.
- Lucchitta, B.K., and Ferguson, H.M., 1988, Ganymede: "Moat" craters compared with palimpsests and basins, *in* Abstracts of papers submitted to the Nineteenth Lunar and Planetary Science Conference, Houston, March 14–18, 1988: Lunar and Planetary Institute, p. 701–702.
- McKinnon, W.B., 1981, Tectonic deformation of Galileo Regio and the limits to the planetary expansion of Ganymede, *in* Lunar and Planetary Science Conference, 12th, Houston, March 16–20, 1981, *Proceedings, part B: Geochimica et Cosmochimica Acta*, p. 1585–1597.
- McKinnon, W.B., and Parmentier, E.M., 1986, Ganymede and Callisto, *in* Burns, J.A., and Matthews, M.S., eds., *Satellites*: Tucson, University of Arizona Press, p. 718–763.
- Murchie, S.L., and Head, J.W., 1988, Possible breakup of dark terrain on Ganymede by large-scale shear faulting: *Journal of Geophysical Research*, v. 93, p. 8795–8824.
- Murchie, S.L., Head, J.W., Helfenstein, Paul, and Plescia, J.B., 1986, Terrain types and local-scale stratigraphy of grooved terrain on Ganymede, *in* Lunar and Planetary Science Conference, 17th, Houston, March 17–21, 1986, *Proceedings, Part 1: Journal of Geophysical Research*, v. 91, suppl., p. E222–E238.
- Murchie, S.L., Head, J.W., and Plescia, J.B., 1989, Crater densities and crater ages of different terrain types on Ganymede: *Icarus*, v. 81, p. 271–297.
- _____, 1990, Tectonic and volcanic evolution of dark terrain and its implications for the internal structure and evolution of Ganymede: *Journal of Geophysical Research*, v. 95, p. 10,743–10,768.
- Parmentier, E.M., and Head, J.W., 1984, A model for the emplacement and solidification of bright terrain on Ganymede, *in* Abstracts of papers submitted to the Fifteenth Lunar and Planetary Science Conference, Houston, March 12–16, 1984: Lunar and Planetary Institute, p. 633–634.
- Passey, Q.R., and Shoemaker, E.M., 1982, Craters and basins on Ganymede and Callisto:


Morphological indicators of crustal evolution, in Morrison, David, ed., *Satellites of Jupiter*: Tucson, University of Arizona Press, p. 379–434.

Schenk, P.M., 1991, Ganymede and Callisto: Complex crater formation and planetary crusts: *Journal of Geophysical Research*, v. 96, p. 15,635–15,664.



QUADRANGLE LOCATION
Number preceded by I refers to published geologic map






INDEX OF Jg-13 MAPPING SOURCES

The rendition of features on this map was controlled by reference to the primary source pictures outlined above. Supplemental source images used during the compilation are listed separately. Copies of various enhancements of these pictures are available from National Space Flight Center, Greenbelt, MD 20771

Primary Source			
VOYAGER 1		VOYAGER 2	
Index No.	Picture No.	Index No.	Picture No.
1	19J1+3	2	401J2-1
		3	404J2-1
		4	407J2-1
		5	413J2-1
		6	431J2-1
		7	494J2-1
		8	506J2-1
		9	518J2-1
		10	521J2-1
		11	530J2-1
		12	573J2-1
		13	662J2-1
		14	666J2-1
		15	684J2-1

Supplemental Source		
VOYAGER 1		VOYAGER 2
Picture No.	Picture No.	Picture No.
0268J1-4	528J2-2	0545J2-2
1207J1-4	0105J2-1	0549J2-1
0015J1+3	0110J2-1	0561J2-1
0593J1-3	0112J2-1	0565J2-1
0318J1-2	0114J2-1	0567J2-1
0788J1-2	0116J2-1	0569J2-1
1651J1-2	0118J2-1	0571J2-1
	0120J2-1	0573J2-1
	0352J2-1	0575J2-1
	0384J2-2	0585J2-1
	0386J2-1	0622J2-1
	0389J2-1	0625J2-1
	0392J2-1	0660J2-1
	0395J2-1	0664J2-1
	0398J2-1	0665J2-1
	0410J2-1	0667J2-1
	0509J2-1	0675J2-1
	0533J2-1	



INDEX OF Jg-12 MAPPING SOURCES

The rendition of features on this map was controlled by reference to the primary source pictures outlined above. Supplemental source images used during the compilation are listed separately. Copies of various enhancements of these pictures are available from National Space Flight Center, Greenbelt, MD 20771

Primary Source			
VOYAGER 2			
Index No.	Picture No.	Index No.	Picture No.
1	528J2-002	21	527J2-001
2	116J2-001	22	530J2-001
3	120J2-001	23	558J2-001
4	124J2-001	24	562J2-001
5	126J2-001	25	564J2-001
6	395J2-001	26	566J2-001
7	407J2-001	27	568J2-001
8	482J2-001	28	570J2-001
9	485J2-001	29	572J2-001
10	488J2-001	30	574J2-001
11	491J2-001	31	576J2-001
12	494J2-001	32	578J2-001
13	497J2-001	33	580J2-001
14	500J2-001	34	582J2-001
15	503J2-001	35	584J2-001
16	509J2-001	36	586J2-001
17	512J2-001	37	588J2-001
18	515J2-001	38	624J2-001
19	521J2-001	39	626J2-001
20	524J2-001		

Supplemental Source	
VOYAGER 2	
Picture No.	Picture No.
105J2-001	549J2-001
122J2-001	564J2-001
352J2-001	567J2-001
376J2-001	575J2-001
384J2-001	585J2-001
470J2-001	625J2-001
500J2-001	665J2-001